



**General Certificate of Education (A-level)  
January 2011**

**Mathematics**

**MPC4**

**(Specification 6360)**

**Pure Core 4**

***Mark Scheme***

---

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: [aqa.org.uk](http://aqa.org.uk)

Copyright © 2011 AQA and its licensors. All rights reserved.

**Copyright**

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

### Key to mark scheme abbreviations

|              |  |
|--------------|--|
| M            | mark is for method   |
| m or dM      | mark is dependent on one or more M marks and is for method         |
| A            | mark is dependent on M or m marks and is for accuracy              |
| B            | mark is independent of M or m marks and is for method and accuracy |
| E            | mark is for explanation  |
| ✓ or ft or F | follow through from previous incorrect result                      |
| CAO          | correct answer only  |
| CSO          | correct solution only  |
| AWFW         | anything which falls within  |
| AWRT         | anything which rounds to   |
| ACF          | any correct form   |
| AG           | answer given   |
| SC           | special case   |
| OE           | or equivalent  |
| A2,1         | 2 or 1 (or 0) accuracy marks                                       |
| -x EE        | deduct $x$ marks for each error                                    |
| NMS          | no method shown  |
| PI           | possibly implied   |
| SCA          | substantially correct approach                                     |
| c            | candidate  |
| sf           | significant figure(s)  |
| dp           | decimal place(s)   |

### No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

**Otherwise we require evidence of a correct method for any marks to be awarded.**

**MPC4**

| Q            | Solution   | Marks | Total    | Comments  |
|--------------|--|-------|----------|---|
| 1(a)         | $R = \sqrt{29}$  | B1    | 3        | Accept 5.4 or 5.38, 5.39, 5.385....<br><br>Condone $\alpha = 68.20^\circ$ |
|              | $R\sin\alpha = 5$ or $R\cos\alpha = 2$ or $\tan\alpha = \frac{5}{2}$ | M1    |          |   |
|              | $\alpha = 68.2^\circ$  | A1    |          |   |
| (b)(i)       | (maximum value =) $\sqrt{29}$  | B1ft  | 1        | ft on $R$   |
| (ii)         | $\sin(x + \alpha) = 1$   | M1    | 2        | Or $x + \alpha = 90$ , $x + \alpha = \frac{\pi}{2}$<br><br>No ISW         |
|              | $x = 21.8^\circ$ only  | A1    |          |   |
| <b>Total</b> |  |       | <b>6</b> |   |



**MPC4 (cont)**

| Q         | Solution  | Marks                               | Total      | Comments |
|-----------|---|-------------------------------------|------------|----------|
| 2(a)(iii) | <p><b>Alternative</b></p> $\frac{f(x) + q(x)}{f(x)}, \text{ where } q \text{ is a quadratic expression}$ $= 1 + \frac{(3x+1)(x+2)}{(3x+1)(3x-1)(x+2)}$ $= 1 + \frac{1}{3x-1}$ | <p>(M1)</p> <p>(A1)</p> <p>(A1)</p> | <p>(3)</p> |          |

## MPC4 (cont)

| Q   | Solution   | Marks | Total | Comments  |  |
|---|--|-------|-------|---|--|
| 3(a)  | $3+9x = A(3+5x) + B(1+x)$  | M1    | 3     | PI by correct A and B   |  |
|   | $x = -1 \quad x = -\frac{3}{5}$  | m1    |       | Substitute two values of $x$ and solve for $A$ and $B$ .                  |  |
|   | $A = 3 \quad B = -6$   | A1    |       |   |  |
|   | <b>Alternative</b><br>Equating coefficients  |       |       |   |  |
|   | $3+9x = A(3+5x) + B(1+x)$  | (M1)  | (3)   | Set up simultaneous equations and solve.<br>Condone 1 error.              |  |
|   | $3 = 3A + B$   | (m1)  |       |   |  |
|   | $9 = 5A + B$   |       |       |   |  |
|   | $A = 3 \quad B = -6$   | (A1)  |       |   |  |
|   | <b>Alternative</b><br>Cover up rule  |       |       |   |  |
|   | $x = -1 \quad A = \frac{3-9}{3-5}$   | (M1)  | (3)   | $x = -1$ and $x = -\frac{3}{5}$<br>and attempt to find $A$ and $B$ .      |  |
| $x = -\frac{3}{5} \quad B = \frac{3-\frac{27}{5}}{1-\frac{3}{5}}$ |  |       |       |   |  |
| $A = 3 \quad B = -6$  | (A1<br>A1)   |       |       |   |  |
| (b)   | $(1+x)^{-1} = 1-x+kx^2$  |       | 7     | SC NMS<br>$A$ and $B$ both correct; 3/3<br>One of $A$ and $B$ correct 1/3 |  |
|   | $= 1-x+x^2$  | M1    |       |   |  |
|   | $(3+5x)^{-1} = 3^{-1}(1+\frac{5}{3}x)^{-1}$  | A1    |       |   |  |
|   | $(1+\frac{5}{3}x)^{-1} = 1-\frac{5}{3}x+(\frac{5}{3}x)^2$                          | B1    |       |   |  |
|   | $= 1-\frac{5}{3}x+\frac{25}{9}x^2$   | M1    |       |   |  |
|   | $\frac{3+9x}{(1+x)(3+5x)}$   | A1    |       |   |  |
|   | $= 3(1-x+x^2) - 6 \times 3^{-1} \left( 1 - \frac{5}{3}x + \frac{25}{9}x^2 \right)$ | M1    |       |   |  |
|   | $= 1 + \frac{1}{3}x - \frac{23}{9}x^2$   | A1    |       |   |  |
|   |  |       |       |   |  |
|   |  |       |       |   |  |

**MPC4 (cont)**

| <b>Q</b> | <b>Solution</b>   | <b>Marks</b> | <b>Total</b> | <b>Comments</b>               |
|----------|---|--------------|--------------|-------------------------------|
| (c)      | $\frac{5x}{3} < 1$ oe or $\frac{5x}{3} > -1$ oe         | M1           |              | Condone $\leq$ instead of $<$ |
|          | $ x  < \frac{3}{5}$ or $-\frac{3}{5} < x < \frac{3}{5}$ | A1           | <b>2</b>     | CAO                           |
|          |   |              | <b>12</b>    |                               |

**MPC4 (cont)**

| Q       | Solution   | Marks                        | Total                 | Comments   |
|---------|--|------------------------------|-----------------------|--|
| 4(a)(i) | $\frac{dx}{dt} = 3e^t$ $\frac{dy}{dt} = 2e^{2t} + 2e^{-2t}$<br><br>$t = 0$ gradient = $\frac{4}{3}$  | M1<br><br>A1<br><br>A1       | <br><br><br>3         | Both derivatives attempted and one correct<br>Both correct<br><br>cso    Condone $\frac{dy}{dx} = \frac{4}{3}$ |
| (ii)    | $y = \frac{4}{3}(x-3)$ oe  | B1ft                         | 1                     | ft on non-zero gradient  |
| (b)     | $e^{2t} = \frac{x^2}{9}$ or $9e^{2t} = x^2$ or $e^t = \frac{x}{3}$ or $e^{2t} = \left(\frac{x}{3}\right)^2$<br><br>or $t = \ln\left(\frac{x}{3}\right)$ or $2t = \ln\left(\frac{x^2}{9}\right)$<br><br>$y = \frac{x^2}{9} - \frac{9}{x^2}$ | <br><br>M1<br><br><br><br>A1 | <br><br><br><br><br>2 | <br><br><br><br><br>Equation required  |
|         |  |                              | <b>6</b>              |  |

**MPC4 (cont)**

| Q    | Solution  | Marks    | Total    | Comments   |
|------|---|----------|----------|--|
| 5(a) | $m = 10 \times 2^{-\frac{14}{8}}$<br>$\approx 3 \text{ (gm)}$                               | M1<br>A1 | 2        | Condone 2.97 or better<br>NOT 2.9 as final answer  |
| (b)  | $2^{-\frac{d}{8}} = \frac{1}{16}$<br>$\frac{d}{8} = 4 \Rightarrow d = 32$                   | M1<br>A1 | 2        | cso  |
| (c)  | $0.01m_0 = m_0 \times 2^{-\frac{t}{8}}$<br>$\ln(0.01) = -\frac{t}{8} \ln(2)$<br>$t = 53.15$ | M1<br>M1 |          | $m_0$ can be numerical<br>Take logs correctly from their equation<br>leading to a linear equation in $t$ . |
|      | $n = 54$  | A1       | 3        | cso  |
|      |   |          | <b>7</b> |  |

| Q  | Solution  | Marks | Total | Comments  |
|--|---|-------|-------|---|
| 6(a)(i)  | $\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$   | B1    | 3     | Condone numerator as $\tan x + \tan x$<br>Multiplying throughout by their denominator<br><b>AG</b> Must show $\tan x = 0$ <b>and</b> $\tan^2 x = 3$ |
|  | $2 \tan x + \tan x(1 - \tan^2 x) = 0$   | M1    |       |   |
|  | $\tan x = 0$  | A1    |       |   |
|  | $\text{or } (2 + 1 - \tan^2 x) = 0 \Rightarrow \tan^2 x = 3$  |       |       |   |
|  | <b>Alternative</b>  |       |       |   |
|  | $\tan 2x = \frac{\sin 2x}{\cos 2x} = \frac{2 \sin x \cos x}{\cos^2 x - \sin^2 x}$   |       |       |   |
|  | $\frac{2 \sin x \cos x}{\cos^2 x - \sin^2 x} + \frac{\sin x}{\cos x} = 0$   | (B1)  |       |   |
|  | $2 \sin x \cos^2 x + \sin x(\cos^2 x - \sin^2 x) = 0$   |       |       |   |
|  | $\sin x(2 \cos^2 x + \cos^2 x - \sin^2 x) = 0$  | (M1)  |       |   |
|  | $\Rightarrow \sin x = 0 \left. \vphantom{\begin{matrix} \Rightarrow \sin x = 0 \\ \Rightarrow \tan x = 0 \end{matrix}} \right\} \text{ and } 3 \cos^2 x = \sin^2 x$ | (A1)  |       |   |
| $\Rightarrow \tan x = 0 \left. \vphantom{\begin{matrix} \Rightarrow \sin x = 0 \\ \Rightarrow \tan x = 0 \end{matrix}} \right\} \text{ and } \tan^2 x = 3$ |   |       |       |   |
| (ii)   | $x = 60$ <b>AND</b> $x = 120$   | B1    | 1     | Condone extra answers outside interval eg 0 and 180   |
| (b)(i)   | $2 \sin x \cos x = \cos x \cdot f(x)$   | M1    | 3     | Where $f(x) = \cos^2 x - \sin^2 x$<br>or $2 \cos^2 x - 1$ or $1 - 2 \sin^2 x$<br><b>AG</b>  |
|  | $2 \sin x \cos x = \cos x(1 - 2 \sin^2 x)$  | A1    |       |   |
|  | $(\cos x \neq 0) \quad 2 \sin x = 1 - 2 \sin^2 x$   | A1    |       |   |
|  | $2 \sin^2 x + 2 \sin x - 1 = 0$   |       |       |   |

|             |   |                            |                       |   |
|-------------|---|----------------------------|-----------------------|---|
| <b>(ii)</b> | $\sin x = \frac{-2 \pm \sqrt{4 - 4 \times 2 \times (-1)}}{2 \times 2}$ $\sin x = \frac{-2 \pm 2\sqrt{3}}{4}$ $\left. \begin{array}{l} \sin x = \frac{-1 - \sqrt{3}}{2} \text{ has no solution} \\ \sin x = \frac{\sqrt{3} - 1}{2} \end{array} \right\}$ | M1<br><br>A1<br><br><br>E1 | <br><br><br><br><br>3 | Correct use of quadratic formula or completing the square or correct factors<br><br>$\sqrt{12}$ must be simplified and must have $\pm$<br><br>Reject one solution and state correct solution. |
|             |   |                            | <b>10</b>             |   |

**MPC4**


| Q           | Solution  | Marks   | Total     | Comments  |                                     |
|-------------|---|---|-----------|---|-------------------------------------|
| 7<br>(a)(i) | $\int \frac{dx}{\sqrt{x}} = \int \sin\left(\frac{t}{2}\right) dt$ | B1  | 3         | Correct separation;<br>condone missing integral signs.              |                                     |
|             | $2\sqrt{x} = -2 \cos\left(\frac{t}{2}\right) (+k)$                | M1  |           | $p\sqrt{x} = q \cos\left(\frac{t}{2}\right)$<br>Condone missing + k |                                     |
|             | $x = \left(-\cos\left(\frac{t}{2}\right) + C\right)^2$            | A1  |           | Must have previous line correct                                     |                                     |
|             | (ii)  | $(1,0) \quad 2 = -2 + k \text{ or } 1 = (-1 + C)^2$   | M1        | 3   | Use (1,0) to find a constant        |
|             |   | $k = 4 \text{ or } C = 2$                             | A1ft      |   | ft on $C = p - q$ from (a)(i)       |
|             |   | $x = \left(2 - \cos\left(\frac{t}{2}\right)\right)^2$ | A1        |   | cso applies to (a)(ii)              |
|             | (b)(i)  | Greatest height when $\cos(bt) = -1$                  | M1        | 2   | ft is (their $a + 1$ ) <sup>2</sup> |
|             |   | Greatest height = 9 (m)                               | A1ft      |   |                                     |
|             | (ii)  | $\cos\left(\frac{t}{2}\right) = 2 - \sqrt{5}$         | M1        | 2   | $\cos bt = a - \sqrt{5}$            |
|             |   | $t = 2 \cos^{-1}(2 - \sqrt{5}) = 3.6$ (seconds 1dp)   | A1        |   | condone 3.6 or better (3.618.....)  |
|             |   |   | <b>10</b> |   |                                     |

## MPC4 (cont)

| Q       | Solution  | Marks | Total     | Comments   |
|---------|---|-------|-----------|--|
| 8(a)(i) | $\overrightarrow{AB} = \begin{bmatrix} 6 \\ 0 \\ 3 \end{bmatrix} - \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix}$   | M1    | 2         | $\pm(\overrightarrow{OB} - \overrightarrow{OA})$ implied by 2 correct components   |
|         |   | A1    |           |  |
| (ii)    | $\begin{bmatrix} 3 \\ 2 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = 6 - 2 - 3 = 1$ $\cos \theta = \frac{sp}{\sqrt{14}\sqrt{14}}$ $\cos \theta = \frac{1}{14} \quad \theta = 85.9^\circ$  | M1    | 4         | Scalar product with correct vectors; allow one component error.<br>ft on $\overrightarrow{AB}$<br><br>Correct form for $\cos \theta$ with one correct modulus<br><br>cso 85.9 or better  |
|         |   | A1ft  |           |  |
|         |   | m1    |           |  |
| (b)(i)  | $\overrightarrow{OD} = \begin{bmatrix} 3 \\ -2 \\ 4 \end{bmatrix} + 2 \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix} = \begin{bmatrix} 7 \\ -4 \\ 10 \end{bmatrix}$   | M1    | 2         | Implied by 2 correct components<br><br>$\mathbf{r} =$ or $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ required ft on $\overrightarrow{AB}$  |
|         |   | A1ft  |           |  |
| (ii)    | $\overrightarrow{BC} = \overrightarrow{OC} - \overrightarrow{OB} = \begin{bmatrix} 1+3p \\ -4+2p \\ 7-p \end{bmatrix}$ $\overrightarrow{AD} = \begin{bmatrix} 4 \\ -2 \\ 6 \end{bmatrix} \quad  \overrightarrow{BC}  = \sqrt{56}$ $(1+3p)^2 + (-4+2p)^2 + (7-p)^2 = 56$ $14p^2 - 24p + 66 = 56$ $7p^2 - 12p + 5 = 0$ $(7p-5)(p-1) = 0$ $p = \frac{5}{7} \text{ and } p = 1$ $C \text{ is at } \left(9\frac{1}{7}, -2\frac{4}{7}, 9\frac{2}{7}\right)$ | M1    | 6         | $\mu = p$ at C<br>Find $\overrightarrow{BC}$ in terms of $p$<br><br>PI B1 is for $ \overrightarrow{BC}  = \sqrt{56}$<br><br>ft on $\overrightarrow{BC}$<br>Simplification to quadratic equation with all terms on one side<br><br>Exact fraction required<br><br>cso Accept as column vector |
|         |   | B1ft  |           |  |
|         |   | m1    |           |  |
|         |   | m1    |           |  |
|         |   | A1    |           |  |
|         |   | A1    |           |  |
|         |   | A1    |           |  |
|         |   |       | <b>14</b> |  |



## MPC4 (cont)

| Q  | Solution  | Marks          | Total     | Comments  |
|--|---|----------------|-----------|---|
| 8(b)(ii)   | Alternative : using symmetry (i)  |                |           |   |
|  | $ \overline{AD}  =  \overline{BC}  = \sqrt{56}$   | (B1ft)         |           | $\overline{AD} = \begin{bmatrix} 4 \\ -2 \\ 6 \end{bmatrix}$  |
|  | $ \overline{DC}  =  \overline{AB}  -  \overline{AD}  \cos \theta -  \overline{BC}  \cos \theta$       | (M1)           |           | Substitute values and evaluate<br>$ \overline{AB}  -  \overline{AD}  \cos \theta -  \overline{BC}  \cos \theta$ |
|  | $ \overline{DC}  = \frac{10}{\sqrt{14}}$  | (A1ft)         |           | F on $\overline{AB}$ and $\cos \theta$  |
|  | $ \overline{DC}  = p  \overline{AB}  \Rightarrow \frac{10}{\sqrt{14}} = p \sqrt{14}$                  | (m1)           |           | Set up equation in $p$  |
|  | $p = \frac{5}{7}$   | (A1)           |           |   |
|  | $C$ is at $\left(9\frac{1}{7}, -2\frac{4}{7}, 9\frac{2}{7}\right)$                                    | (A1)           | (6)       |                             |
|  | Alternative using symmetry (ii)   |                |           |   |
|  | $ \overline{AD}  = \sqrt{56}$   | (B1ft)         |           |   |
|  | $ \overline{AE}  =  \overline{AD}  \cos \theta = \sqrt{56} \times \frac{1}{14} = \frac{2}{\sqrt{14}}$ | (M1)<br>(A1ft) |           | Substitute values and evaluate<br>for $ \overline{AD}  \cos \theta$ . F on $\cos \theta$                        |
|  | $ \overline{AE}  = q  \overline{AB}  \Rightarrow \frac{2}{\sqrt{14}} = q \sqrt{14}$                   | (m1)           |           | Set up equation to find $p$   |
|  | and $ \overline{AE}  =  \overline{FB}  \Rightarrow p = 1 - 2q$  |                |           |   |
| $q = \frac{2}{14} \quad p = \frac{5}{7}$                           | (A1)  |                |           |   |
| $C$ is at $\left(9\frac{1}{7}, -2\frac{4}{7}, 9\frac{2}{7}\right)$ | (A1)  | (6)            |           |   |
|  | <b>TOTAL</b>  |                | <b>75</b> |   |