

General Certificate of Education  
June 2009  
Advanced Level Examination



**MATHEMATICS**  
**Unit Pure Core 3**

**MPC3**

Friday 5 June 2009 1.30 pm to 3.00 pm

**For this paper you must have:**

- an 8-page answer book
- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Time allowed: 1 hour 30 minutes

**Instructions**

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MPC3.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.

**Information**

- The maximum mark for this paper is 75.
- The marks for questions are shown in brackets.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

---

Answer **all** questions.

---

1 (a) The curve with equation

$$y = \frac{\cos x}{2x + 1}, \quad x > -\frac{1}{2}$$

intersects the line  $y = \frac{1}{2}$  at the point where  $x = \alpha$ .

(i) Show that  $\alpha$  lies between 0 and  $\frac{\pi}{2}$ . (2 marks)

(ii) Show that the equation  $\frac{\cos x}{2x + 1} = \frac{1}{2}$  can be rearranged into the form

$$x = \cos x - \frac{1}{2} \quad (1 \text{ mark})$$

(iii) Use the iteration  $x_{n+1} = \cos x_n - \frac{1}{2}$  with  $x_1 = 0$  to find  $x_3$ , giving your answer to three decimal places. (2 marks)

(b) (i) Given that  $y = \frac{\cos x}{2x + 1}$ , use the quotient rule to find an expression for  $\frac{dy}{dx}$ . (3 marks)

(ii) Hence find the gradient of the normal to the curve  $y = \frac{\cos x}{2x + 1}$  at the point on the curve where  $x = 0$ . (2 marks)

2 The functions  $f$  and  $g$  are defined with their respective domains by

$$f(x) = \sqrt{2x + 5}, \quad \text{for real values of } x, \quad x \geq -2.5$$
$$g(x) = \frac{1}{4x + 1}, \quad \text{for real values of } x, \quad x \neq -0.25$$

(a) Find the range of  $f$ . (2 marks)

(b) The inverse of  $f$  is  $f^{-1}$ .

(i) Find  $f^{-1}(x)$ . (3 marks)

(ii) State the domain of  $f^{-1}$ . (1 mark)

(c) The composite function  $fg$  is denoted by  $h$ .

(i) Find an expression for  $h(x)$ . (1 mark)

(ii) Solve the equation  $h(x) = 3$ . (3 marks)

3 (a) Solve the equation  $\tan x = -\frac{1}{3}$ , giving all the values of  $x$  in the interval  $0 < x < 2\pi$  in radians to two decimal places. (3 marks)

(b) Show that the equation

$$3 \sec^2 x = 5(\tan x + 1)$$

can be written in the form  $3 \tan^2 x - 5 \tan x - 2 = 0$ . (1 mark)

(c) Hence, or otherwise, solve the equation

$$3 \sec^2 x = 5(\tan x + 1)$$

giving all the values of  $x$  in the interval  $0 < x < 2\pi$  in radians to two decimal places. (4 marks)


Turn over ►

- 
- 4 (a) Sketch the graph of  $y = |50 - x^2|$ , indicating the coordinates of the point where the graph crosses the  $y$ -axis. *(3 marks)*
- (b) Solve the equation  $|50 - x^2| = 14$ . *(3 marks)*
- (c) Hence, or otherwise, solve the inequality  $|50 - x^2| > 14$ . *(2 marks)*
- (d) Describe a sequence of two geometrical transformations that maps the graph of  $y = x^2$  onto the graph of  $y = 50 - x^2$ . *(4 marks)*
- 5 (a) Given that  $2 \ln x = 5$ , find the exact value of  $x$ . *(1 mark)*
- (b) Solve the equation


$$2 \ln x + \frac{15}{\ln x} = 11$$

giving your answers as exact values of  $x$ . *(5 marks)*

- 6 The diagram shows the curve with equation  $y = \sqrt{100 - 4x^2}$ , where  $x \geq 0$ .



- (a) Calculate the volume of the solid generated when the region bounded by the curve shown above and the coordinate axes is rotated through  $360^\circ$  about the **y-axis**, giving your answer in terms of  $\pi$ . (5 marks)
- (b) Use the mid-ordinate rule with five strips of equal width to find an estimate for  $\int_0^5 \sqrt{100 - 4x^2} dx$ , giving your answer to three significant figures. (4 marks)
- (c) The point  $P$  on the curve has coordinates  $(3, 8)$ .
- (i) Find the gradient of the curve  $y = \sqrt{100 - 4x^2}$  at the point  $P$ . (3 marks)
- (ii) Hence show that the equation of the tangent to the curve at the point  $P$  can be written as  $2y + 3x = 25$ . (2 marks)
- (d) The shaded regions on the diagram below are bounded by the curve, the tangent at  $P$  and the coordinate axes.



Use your answers to part (b) and part (c)(ii) to find an approximate value for the **total** area of the shaded regions. Give your answer to three significant figures. (5 marks)

Turn over ►

7 (a) Use integration by parts to find  $\int (t - 1) \ln t \, dt$ . *(4 marks)*

(b) Use the substitution  $t = 2x + 1$  to show that  $\int 4x \ln(2x + 1) \, dx$  can be written as  $\int (t - 1) \ln t \, dt$ . *(3 marks)*

(c) Hence find the exact value of  $\int_0^1 4x \ln(2x + 1) \, dx$ . *(3 marks)*

**END OF QUESTIONS**

**There are no questions printed on this page**

**There are no questions printed on this page**